

MEDICAL SCIENCE

To Cite:

Ahangari N, Fatollahzadeh S, Doos Ali Vand H, Sarani Yaztappeh J. Effect of Internet-based versus face-to-face Cognitive Behavioral Therapy for Dependent Personality Disorder: A randomized clinical trial. *Medical Science* 2023; 27: e354ms3223
doi: <https://doi.org/10.54905/disssi.v27i139.e354ms3223>

Authors' Affiliation:

¹PhD Candidate in Clinical Psychology, Department of Clinical Psychology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

²Master of Science, Department of Clinical Psychology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

³Assistant Professor of Clinical Psychology, Department of Clinical Psychology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

*Corresponding author

Assistant Professor of Clinical Psychology, Department of Clinical Psychology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran,

Iran

Email: doosalivand.h@sbmu.ac.ir

Contact List

Nazanin Ahangari
Saina Fatollahzadeh
Hoda Doos Ali Vand
Jafar Sarani Yaztappeh

nazaninahangari@gmail.com
sainafatollahzadeh@gmail.com
doosalivand.h@sbmu.ac.ir
Email: sarani.jafar@yahoo.com

ORCID List

| | |
|------------------------|---------------------|
| Nazanin Ahangari | 0000-0003-2164-1555 |
| Saina Fatollahzadeh | 0000-0001-8303-0667 |
| Hoda Doos Ali Vand | 0000-0002-1814-979X |
| Jafar Sarani Yaztappeh | 0000-0002-6430-202X |

Peer-Review History

Received: 02 July 2023

Reviewed & Revised: 06/July/2023 to 12/September/2023

Accepted: 16 September 2023

Published: 21 September 2023

Peer-review Method

External peer-review was done through double-blind method.

Medical Science

pISSN 2321-7359; eISSN 2321-7367



© The Author(s) 2023. Open Access. This article is licensed under a [Creative Commons Attribution License 4.0 \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

Effect of Internet-based versus face-to-face Cognitive Behavioral Therapy for Dependent Personality Disorder: A randomized clinical trial

Nazanin Ahangari¹, Saina Fatollahzadeh², Hoda Doos Ali Vand^{3*}, Jafar Sarani Yaztappeh¹

ABSTRACT

Background: Dependent Personality Disorder (DPD) is the most prevalent personality disorder, which is associated with several negative consequences. **Objectives:** The current study aimed to investigate the effect of ICBT on DPD and compare it with face-to-face CBT in a sample of DPD patients. **Method:** The participants were adults with DPD recruited from various counseling and psychological clinics in Tehran, Iran. They were randomly assigned to receive face-to-face CBT ($n = 20$) or ICBT ($n = 20$). Depression-Anxiety-Stress Scale-21 (DASS-21) and Interpersonal Problems Questionnaire (IIP-32) were completed at pre- and post-treatment. **Result:** Analysis of Covariance (ANCOVA) showed that the face-to-face CBT group had significantly lower interpersonal problem scores than the ICBT group ($F=23.81$, $p<0.05$). However, no significant difference was observed between the face-to-face CBT and ICBT groups in anxiety ($F=3.68$, $p<0.05$) and depression ($F=1.70$, $p<0.05$). **Conclusion:** The findings support the effects of both ICBT and face-to-face CBT in treating patients with DPD. While both ICBT and face-to-face CBT demonstrated favorable outcomes, our study suggests an advantage of face-to-face CBT in addressing interpersonal problems among these patients. Further research is needed to investigate the long-term effects of ICBT to enhance traditional approaches for DPD.

Keywords: Cognitive Behavioral Therapy, Dependent Personality Disorder, Interpersonal Problems, Anxiety, Depression

1. INTRODUCTION

Dependent Personality Disorder (DPD) is defined as a maladaptive pattern of submissiveness, fear of separation, and clinging behavior. This disorder

typically begins in early adulthood and is characterized by a strong desire to be taken care of and a belief that one is incapable of functioning independently (Association, 2013). Patients with DPD have difficulties making decisions, expressing disagreement, and taking responsibility; they also suffer from passivity, lack of self-confidence, and imbalanced or distorted relationships (Association, 2013). DPD affects 0.49% of the general adult population. Also, some studies have reported 1-2% of DPD prevalence (Anderson et al., 2017; Trull et al., 2010). A previous study reported that the prevalence of DPD was 1.79% in a sample of Iranian college students (Ghaderi et al., 2016). Women are 40% more diagnosed with DPD than men (Crego and Widiger, 2019). Even though DPD is more prevalent than other personality disorders, only a few clinical trials have investigated the effectiveness of therapeutic approaches in these patients (Faith, 2009).

Patients with DPD are more vulnerable to somatization, major depression, panic, and eating disorders than other personality disorders (Barzega et al., 2001; Coyne and Whiffen, 1995; Kemp et al., 2013; Overholser, 1996; Rost et al., 1992; Tisdale et al., 1990). Moreover, DPD is associated with gastrointestinal problems and sleep disturbances (Sachse et al., 2013). There are various treatment approaches for DPD, including psychodynamic and Cognitive Behavioral Therapy (CBT) (Eskedal and Demetri, 2006; Faith, 2009; Sperry, 2013). Time-limited psychodynamic therapy is considered the most suitable approach for individuals diagnosed with DPD. However, it has been found that patients with limited ego strength or significant separation anxiety may not benefit as much from short or long-term psychodynamic therapy compared to alternative treatment options (Eskedal and Demetri, 2006). CBT intends to increase the patient's autonomy and self-efficacy Sperry, (2013), and it is appropriate for addressing the longstanding problems of patients with personality disorders (Beck et al., 2015; Linehan, 1993).

Despite considerable advances in developing, assessing, and disseminating empirically supported treatments for Axis I disorders, only slow progress has been observed for most personality disorders. Most randomized clinical trials for personality disorders are limited to borderline and avoidant personality disorders (Matusiewicz et al., 2010). As a more recent form of CBT, Internet-based Cognitive Behavioral Therapy (ICBT) has growing empirical support for various psychiatric conditions (Andrews et al., 2010; Hedman et al., 2012). Owing to the exponential growth of internet access, ICBT was developed and welcomed by therapists (Webb et al., 2017). The advantages of ICBT over face-to-face CBT are a lower dropout of elders, high fidelity, greater accessibility, greater convenience, reduced cost to patients, and limited professional support for its administration (Andersson and Cuijpers, 2009; Edmonds et al., 2018; Kenter et al., 2015). A systematic review indicated that ICBT is as effective as CBT for mental disorders such as depression and anxiety (Hedman et al., 2012).

However, as personality disorders are complex and require more interaction between patient and therapist, there has been little development of internet-based therapies for these disorders (van-der-Boom et al., 2022). Mahoney et al., (2021) investigated the effectiveness of ICBT for depression and anxiety in patients with comorbid personality disorders. The results showed that comorbid personality disorders were not significantly associated with poorer treatment adherence, higher post-treatment symptom severity, or psychological distress (Mahoney et al., 2021). Most previous studies of internet-based therapy for personality disorders were developed for borderline personality disorders van-der-Boom et al., (2022) and were based on schema therapy and dialectical behavioral therapy (Austin et al., 2020; Jacob et al., 2018; Klein et al., 2018; Klein et al., 2021; Rizvi et al., 2011; van-der-Boom et al., 2022).

There is a clear need for further development and evaluation of CBT effects on understudied personality disorders such as DPD (Matusiewicz et al., 2010). As far as the authors know, no published study has investigated the effects of ICBT on DPD. In the same way, no published study has directly compared ICBT to face-to-face CBT in patients with DPD. Thus, the aim of the current study was to investigate the effect of ICBT on DPD and compare it with face-to-face CBT in a sample of DPD patients.

2. MATERIALS AND METHODS

Participants

Participants were recruited from various counseling and psychological clinics in Tehran, Iran. These patients were self-referred, with a mean age of 38.8 (SD = 6.62). Eligible participants were aged 18 and 65, had a primary diagnosis of DPD, and were fluent in Persian. They were excluded if they met the criteria for a current diagnosis of schizophrenia spectrum disorder, bipolar I disorder, neurocognitive disorder, or substance-related disorder, according to the DSM-5. They were also excluded if they had a diagnosis of antisocial, schizotypal, schizoid, or borderline personality disorder, were at high risk of suicide, had a history of CBT within the last four years, underwent other psychological treatment during the study, or started, stopped, or changed medication doses during the course of the study. Figure 1 depicts the trial design in the Consolidated Standards of Reporting Trials (CONSORT) flowchart.




Figure 1 CONSORT flowchart of participant recruitment, randomization, and retention

Measures

Depression Anxiety Stress Scale-21 (DASS-21)

This 21-item scale Henry and Crawford, (2005) is a brief version of DASS-42. This self-report scale measures emotional states of depression, anxiety, and stress. Responses are rated on a 4-point Likert scale ranging from 0 (Did not apply to me at all) to 3 (Applied to me very much or most of the time). Internal consistency of DASS-21 in a non-clinical sample has been reported to be 0.88, 0.82, 0.90, and 0.93 for depression, anxiety, and stress subscales and the total scale, respectively (Henry and Crawford, 2005). The Persian version of the DASS-21 was found to have an internal consistency of 0.94 in a non-clinical sample (Asghari et al., 2008). In this study, the *Cronbach's* alpha for the depression and anxiety subscales was 0.79 and 0.74, respectively.

Inventory of Interpersonal Problems (IIP-32)

IIP-32 assesses people's difficulties in their relationships with others. This 32-item scale Barkham et al., (1996) is a brief version of the Interpersonal Problems Inventory. The answers are rated on a scale from 0 (Not at all) to 4 (Very much). Moreover, two interpersonal dimensions of agency and communion are calculated based on weighted combinations of these eight subscales (Gómez-Penedo et al., 2022). The subscales of the IIP-32 have exhibited satisfactory levels of internal consistency in both outpatient and non-clinical populations (Barkham et al., 1996). The Persian version of the IIP-32 includes 6 subscales (Assertiveness and sociability, openness, caring, aggression, supportiveness and involvement, and dependency). The IIP-32 Persian version had a *Cronbach's* alpha of 0.82 in an Iranian sample (Fath et al., 2013).

Structured Clinical Interview for DSM-5-Research Version (SCID-5-RV)

SCID-5-RV (First et al., 2015) is a guide for semi-structured interviews for the major DSM-5 diagnoses. The task is carried out by a skilled clinician or healthcare professional who is knowledgeable about the diagnostic criteria and classification of disorders in DSM-5 (First et al., 2015). The SCID-5-RV is generally administered in a single 45-90-minute session (Jiang et al., 2021). The Research Version contains more disorders than the Clinician Version (Blackmore et al., 2022). Several studies have shown that SCID-5-RV has acceptable reliability and validity (First et al., 2015). The Persian version of the SCID-5-RV indicated an acceptable range for internal consistency (0.95-0.99), test-retest reliability (0.60-0.79), and kappa reliability (0.57-0.72) (Mohammadkhani et al., 2020).

Structured Clinical Interview for DSM-5 Personality Disorders (SCID-5-PD)

SCID-5-PD First et al., (2016) is a semi-structured diagnostic interview to assess DSM-5 personality disorders under three clusters of A, B, C, and other specific personality disorders. This semi-structured interview consists of 119 items. The SCID-5-PD can help diagnose personality disorder, either categorically (present or absent) or dimensionally summing the ratings (0, 1, or 2) (First et al., 2016; Somma et al., 2020). SCID-5-PD has been shown to have good interrater reliability in a European sample (Somma et al., 2020). The Persian version of the SCID-5-PD indicated good interrater reliability for all personality disorders except for the schizotypal personality disorder. The kappa of the Persian version of the SCID-5-PD for all personality disorders was higher than 0.4, except for DPD and avoidant personality disorder (Gharraee et al., 2021).

Structured Clinical Interview for DSM-5 Screening Personality Questionnaire (SCID-5-SPQ)

SCID-5-SPQ First et al., (2016) is a self-report screening tool to assess the diagnostic criteria of 10 DSM-5 personality disorders. This tool serves as a preliminary data collection tool before the interview. It contains 106 true-false self-report items (Fjermestad-Noll et al., 2020). The internal consistency of the SCID-5-SPQ Persian version has been reported to be 0.93 (Gharraee et al., 2022).

Procedure and study design

This randomized controlled trial compared ICBT and face-to-face CBT in a sample of adults with DPD. The study was conducted between June and August 2023. All participants provided written informed consent. They were diagnosed with DPD using SCID-5-PD and SCID-5-SPQ. Independent evaluators (psychologists) who were blind to the treatment condition administered SCID-5-PD at baseline. In addition, the participants were asked to complete SCID-5-SPQ. Eligible participants were randomized to face-to-face CBT or ICBT groups. Randomization was achieved by even/odd serial numbers. Patients completed self-report measures (DASS-21 and IIP-32) pre-and post-treatment. The kappa of SCID-5-PD for DPD was below 0.4. The outcome measures were DASS-21 and IIP-32.

Therapists

Therapists were two doctoral students with 4-5 years of experience. They had expertise in treating personality disorders both face-to-face and digitally. They received supervision on request at the clinic's own scheduled supervision hours.

Interventions

The treatment was adapted from the *cognitive behavioral therapy of DSM-5 personality disorders* (Sperry and Sperry, 2015). Face-to-face CBT and ICBT consisted of 15 sessions. Each treatment session lasted 45 minutes once a week.

Face-to-face cognitive behavioral therapy

CBT is based on the assumption that dependency is related to individuals' maladaptive beliefs and related thought processes (Beck et al., 2015). After building a collaborative interaction between patient and therapist, CBT applies specific cognitive and behavioral techniques such as Socratic dialogue, cognitive awareness training, pinpointing and challenging automatic beliefs, assertive communication training, and problem-solving training (Sperry and Sperry, 2015).

Internet-based cognitive behavioral therapy

The content of ICBT sessions was similar to face-to-face CBT, except that the sessions were provided through the native Skyroom platform. This platform does not require the installation of any applications, and users can access the desired event with just one click.

Statistical analyses

Analyses were conducted using SPSS 28 software. The Chi-square and independent t-test analyzed the participants' demographic and clinical characteristics. Changes in the mean scores for depression, anxiety, and interpersonal problems from pre- to post-treatment within the face-to-face CBT and ICBT groups were analyzed with paired t-tests. The Analysis of Covariance (ANCOVA) examined differences in interpersonal problems, depression, and anxiety between the treatment groups. $P < 0.05$ was considered statistically significant.

3. RESULTS

Table 1 summarizes the participants' demographic characteristics. To indicate the normality of the data, we used the skewness and kurtosis and Shapiro-Wilk test. All variables approximated normal distribution (skewness and kurtosis from -2 to +2; $p < 0.05$ on the Shapiro-Wilk test). No significant difference was found in age ($t = 0.55$, $p > 0.05$), gender ($X^2 = 0.75$, $p > 0.05$), marital status ($X^2 = 0.33$, $p > 0.05$), education ($X^2 = 0.88$, $p > 0.05$), occupational status ($X^2 = 0.70$, $p > 0.05$), current medication use ($X^2 = 0.76$, $p > 0.05$), Axis I disorders ($X^2 = 0.543$, $p > 0.05$), and Axis II disorders ($X^2 = 0.124$, $p > 0.05$) between the face-to-face CBT and ICBT groups. Figure 2 and Figure 3 present clinical characteristics of participants.

Table 1 Demographic characteristics of participants

| Variable | | Face-to-face CBT N (%) | ICBT N (%) | Statistical significance |
|---------------------|-------------------|------------------------|------------|--------------------------|
| Age | Mean | 39.40 | 38.20 | .556 |
| | SD | 6.75 | 6.50 | |
| Gender | Women | 14 (66.7) | 13 (59.1) | .755 |
| | Men | 7 (33.3) | 9 (40.9) | |
| Married status | Marital | 16 (76.2) | 13 (59.1) | .332 |
| | Single | 5 (23.8) | 9 (40.9) | |
| Education | High school | 7 (33.3) | 6 (27.3) | .889 |
| | Bachelor's degree | 6 (28.6) | 8 (36.4) | |
| | Master's degree | 5 (23.8) | 6 (27.3) | |
| | Ph.D. degree | 3 (14.3) | 2 (9.1) | |
| Occupational status | Employed | 14 (66.7) | 17 (77.3) | .701 |
| | Unemployed | 5 (23.8) | 4 (18.2) | |
| | Retired | 2 (9.5) | 1 (4.5) | |




Figure 2 Frequency of current medication use in face-to-face CBT and ICBT groups




Figure 3 Frequency of axis I and axis II disorders in face-to-face CBT and ICBT groups

Table 2 shows the means, standard deviations, and paired t-test results for the face-to-face CBT and ICBT groups on all outcome measures at pre- and post-treatment. Based on the paired t-test, the ICBT group significantly improved from pre- to post-treatment in depression ($t(1.41) = 9.99, p < 0.05$), anxiety ($t(1.41) = 12.35, p < 0.05$), and interpersonal problems ($t(1.41) = 12.54, p < 0.05$). Also, a significant change was observed in depression ($t(1.41) = 8.66, p < 0.05$), anxiety ($t(1.41) = 9.75, p < 0.05$), and interpersonal problems ($t(1.41) = 14.56, p < 0.05$) in the face-to-face group.

Table 2 Means, standard deviations, and t-test results for face-to-face and ICBT groups on all outcome measures at pre- and post-treatment

| Variable | Group | Pre-test | Post-test | Mean difference | t value | p-value |
|---------------------------------|--------------|---------------|------------|-----------------|---------|---------|
| Anxiety (DASS-21) | Face-to-face | 10.84a ±2.25b | 7.86±2.23 | 2.99 ±.306c | 9.75 | < .001 |
| | ICBT | 11.50±2.20 | 8.95±1.92 | 2.55±.207 | 12.35 | < .001 |
| Depression (DASS-21) | Face-to-face | 14.67±2.70 | 10.75±2.32 | 3.91±.452 | 8.66 | < .001 |
| | ICBT | 15.31±2.96 | 11.63±2.54 | 3.68±.368 | 9.99 | < .001 |
| Assertiveness and sociability | Face-to-face | 15.05±2.82 | 12.14±2.69 | 2.91±.351 | 8.27 | < .001 |
| | ICBT | 15.82±2.74 | 14.00±2.27 | 1.82±.340 | 5.35 | < .001 |
| Openness | Face-to-face | 13.71±2.80 | 9.41±2.17 | 4.30±.592 | 7.28 | < .001 |
| | ICBT | 12.86±2.24 | 10.87±1.84 | 1.99±1.127 | 4.79 | < .001 |
| Caring | Face-to-face | 14.83±3.73 | 11.58±1.67 | 3.25±.689 | 4.72 | < .001 |
| | ICBT | 14.13±2.86 | 11.05±1.34 | 3.07±.415 | 7.40 | < .001 |
| Aggression | Face-to-face | 7.95±1.66 | 7.29±1.45 | .67±.354 | 1.88 | .074 |
| | ICBT | 7.59±1.71 | 7.27±1.32 | .32±.250 | 1.28 | .216 |
| Supportiveness and involvement | Face-to-face | 12.81±3.16 | 12.05±2.59 | .76±.447 | 1.71 | .104 |
| | ICBT | 12.50±3.50 | 12.23±3.15 | .27±.324 | .84 | .409 |
| Dependency | Face-to-face | 9.71±1.91 | 9.18±1.41 | 3.43±.486 | 7.06 | < .001 |
| | ICBT | 6.29±1.53 | 9.18±1.46 | 2.18±.252 | 8.67 | < .001 |
| Interpersonal problems (IIP-32) | Face-to-face | 74.07±8.34 | 58.75±8.24 | 15.31±1.05 | 14.56 | < .001 |
| | ICBT | 72.08±9.19 | 62.43±7.25 | 9.66±.770 | 12.54 | < .001 |

a = Mean, b = Standard deviation, c = Standard error of mean

The ANCOVA showed that the face-to-face CBT group had significantly lower interpersonal problem scores (mean=58.11, SE=0.719) than the ICBT group (mean=63.04, SE=0.702) ($F=23.81, df=1,41, p < 0.05$). However, no significant difference was established

between the face-to-face CBT and ICBT groups in anxiety ($F=3.68$, $df =1,41$, $p>0.05$) and depression ($F=1.70$, $df =1,41$, $p>0.05$) (Table 3).

Table 3 Comparison of mean scores of all outcome measures in face-to-face CBT and ICBT groups

| Variable | Adjusted mean | | Mean difference | F value | p-value | η^2 |
|--------------------------------|-------------------|------------------|------------------|---------|---------|----------|
| | ICBT | Face-to-face | | | | |
| Anxiety | 8.08a \pm .226b | 8.73 \pm .221 | .649 \pm .338 | 3.68 | .062 | .07 |
| Depression | 10.92 \pm .297 | 11.47 \pm .290 | .541 \pm .416 | 1.70 | .200 | .04 |
| Assertiveness and sociability | 12.43 \pm .314 | 13.72 \pm .308 | 1.29 \pm .443 | 8.52 | .006 | .18 |
| Openness | 9.17 \pm .341 | 11.10 \pm .333 | 1.92 \pm .478 | 16.22 | < .001 | .29 |
| Caring | 11.42 \pm .434 | 11.21 \pm .424 | .210 \pm .609 | .12 | .732 | < .01 |
| Aggression | 7.18 \pm .265 | 7.37 \pm .259 | .188 \pm .371 | .26 | .615 | .01 |
| Supportiveness and involvement | 11.93 \pm .352 | 12.34 \pm .343 | .412 \pm .492 | .70 | .407 | .02 |
| Dependency | 6.16 \pm .314 | 7.12 \pm .308 | .964 \pm .443 | 4.73 | .036 | .11 |
| Interpersonal problems | 58.11 \pm .719 | 63.04 \pm .702 | 4.93 \pm 1.010 | 23.81 | < .001 | .37 |

a = Mean, b = Standard error of mean

4. DISCUSSION

This study investigated the effect of ICBT on DPD and compared it with face-to-face CBT in a sample of DPD patients. Our findings indicated that ICBT was similar to face-to-face CBT in reducing depression and anxiety symptoms in patients with DPD. However, face-to-face CBT was more effective than ICBT in improving interpersonal problems. The primary finding of the current study was that ICBT was as effective as face-to-face CBT in reducing depression and anxiety symptoms in DPD patients. This finding aligns with previous studies showing comparable effectiveness of face-to-face and internet-based treatment for depression and anxiety (Komariah et al., 2022; Lappalainen et al., 2014; Wagner et al., 2014). However, a study by Wagner et al., (2014) showed that internet-based treatment, compared to face-to-face treatment, had a sustained symptom reduction, suggesting potential long-term effectiveness that may be unique to internet-based interventions (Wagner et al., 2014).

Our study did not include follow-up assessments, so the longitudinal effects of face-to-face CBT compared to ICBT are unclear. Additionally, Casey et al., (2014) found that patients with depression do not prefer internet-based therapy due to lower perceived barriers to treatment (e.g., time constraints and participation restrictions) (Casey et al., 2014). Therefore, our findings regarding the equal effectiveness of ICBT and face-to-face CBT for depression and anxiety symptoms are consistent with the notion that perceived barriers to treatment may not significantly impact treatment outcomes. Furthermore, it is important to note that face-to-face CBT and ICBT share similar techniques, processes, and treatment goals, which could explain the comparable effectiveness of both approaches for depression and anxiety symptoms. The second finding of the current study was that face-to-face CBT and ICBT effectively improved interpersonal problems.

However, face-to-face CBT was more effective than ICBT in reducing interpersonal problems. This finding aligns with previous studies supporting CBT as a well-established and effective treatment for Personality Disorders (PDs), including DPD (Matusiewicz et al., 2010; Zanin and Valerio, 2004). More specifically, the intervention approach employed in this study encompassed various components, including assertiveness training and problem-solving skills, which have been previously established as effective techniques for addressing and ameliorating interpersonal problems (Spence et al., 1994; Sperry and Sperry, 2015; Tisdelle and Lawrence, 1986). In addition, CBT has been found to place a greater emphasis on addressing maladaptive schemas in treating PDs than Axis 1 disorders (Rafaeli, 2009). Beck and colleagues linked dependency to a maladaptive incompetence schema (Gude et al., 2004). Our treatment approach concentrated on the maladaptive schemas of dependency/incompetence and failure to achieve in individuals with DPD (Sperry and Sperry, 2015).

Addressing and modifying these maladaptive schemas could notably impact behavioral patterns and subsequent improvement of interpersonal problems. Previous studies highlight the importance of the therapeutic relationship in treating PDs (Muran et al., 2018). Several studies have reported lower alliance levels in internet-based therapy compared to face-to-face therapy Kiropoulos et al., (2008), Weineland et al., (2020), which may explain the difference in the effectiveness of face-to-face CBT compared to ICBT for

DPD patients. However, it is worth noting that other studies have found similar levels of alliance in both modalities (Kaiser et al., 2021). Therefore, it is important to investigate the underlying factors influencing alliance in different therapeutic settings. In general, while the effectiveness of face-to-face CBT and ICBT was comparable for depression and anxiety symptoms, individuals with DPD may be particularly sensitive to the quality of alliance.

Overall, face-to-face CBT remains an effective treatment for DPD, but ICBT may be a viable option for patients who cannot access in-person therapy. Mental health professionals must consider individual patient needs and preferences when selecting treatment options for DPD. Despite the promising findings of the current study, some limitations warrant mention. First, the kappa of the Persian version of the SCID-5-PD for DPD has been reported to be low. However, we used the SCID-5-SPQ in addition to SCID-5-PD to acknowledge the diagnosis of DPD. Second, the current study did not include a follow-up assessment. Thus, longer-term associations between treatment and outcome measures are unknown.

Future studies should consider follow-up assessments. Third, since ICBT studies for DPD are scarce, comparing our results with previous studies was difficult. More studies are needed to investigate the effect of ICBT in DPD patients. Finally, as the present study did not investigate the mechanisms of change, no firm conclusions can be drawn about therapy processes. Future studies should more specifically understand how these outcomes are achieved and whether there are differences in mechanisms across treatments.

5. CONCLUSION

This study investigated the effect of ICBT on DPD and compared it with face-to-face CBT in a sample of DPD patients. Our findings indicated that both ICBT and face-to-face CBT effectively reduced depression and anxiety, while face-to-face CBT showed an advantage in decreasing interpersonal problems. However, it is important to note that ICBT may still have potential as an alternative treatment option, particularly in remote areas where access to CBT may be limited. Further research is needed to explore the long-term effects of ICBT and its potential to complement or enhance traditional CBT approaches.

Acknowledgment

The authors thank those who participated in this study.

Author contributions

NA and HD: Conceptualization, design, and methodology. JS: Data collection and formal analysis. HD: Supervision. SF, JS, and NA: Investigation and project administration. NA and SF: Writing the original draft. HD: Revising the manuscript.

Ethical approval

The Medical Ethics Committee of Shahid Beheshti University of Medical Sciences approved the study (Ethical approval code: IR.SBMU.MSP.REC.1402.060).

Informed consent

Written and oral informed consent was obtained from all individual participants included in the study.

Funding

This study was funded by the School of Medicine, Shahid Beheshti University of Medical Sciences (grant number 43002323).

Conflict of interest

The authors declare that there is no conflict of interests.

Data and materials availability

All data sets collected during this study are available upon reasonable request from the corresponding author.

REFERENCES AND NOTES

1. Anderson D, Lord C, Risi S, Di-Lavore P, Shulman C, Thurm A, Pickles A. American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders. Washington, DC: Author. The Linguistic and Cognitive Effects of Bilingualism on Children with Autism Spectrum

- Disorders 2017; 21(1):1-197. doi: 10.1176/appi.books.9780890425596
2. Andersson G, Cuijpers P. Internet-based and other computerized psychological treatments for adult depression: a meta-analysis. *Cogn Behav Ther* 2009; 38(4):196-205. doi: 10.1080/16506070903318960
3. Andrews G, Cuijpers P, Craske MG, McEvoy P, Titov N. Computer therapy for the anxiety and depressive disorders is effective, acceptable and practical health care: a meta-analysis. *PloS One* 2010; 5(10):e13196. doi: 10.1371/journal.pone.0013196
4. Asghari A, Saed F, Dibajnia P. Psychometric properties of the Depression Anxiety Stress Scales-21 (DASS-21) in a non-clinical Iranian sample. *Int J psychol* 2008; 2(2):82-102.
5. Association AP. Diagnostic and Statistical Manual of Mental Disorders (DSM-V). 5th Edition, American Psychiatric Publishing, Washington DC, 2013. doi: 10.1176/appi.books.9780890425596
6. Austin SF, Jansen JE, Petersen CJ, Jensen R, Simonsen E. Mobile app integration into dialectical behavior therapy for persons with borderline personality disorder: qualitative and quantitative study. *JMIR Ment Health* 2020; 7(6):e14913. doi: 10.2196/14913
7. Barkham M, Hardy GE, Startup M. The IIP-32: A short version of the Inventory of Interpersonal Problems. *Br J Clin Psychol* 1996; 35(1):21-35. doi: 10.1111/j.2044-8260.1996.tb01159.x PMID:8673033
8. Barzega G, Maina G, Venturello S, Bogetto F. Gender-related distribution of personality disorders in a sample of patients with panic disorder. *Eur Psychiatry* 2001; 16(3):173-9. doi: 10.1016/S0924-9338(01)00560-0
9. Beck AT, Davis DD, Freeman A. Cognitive therapy of personality disorders: Guilford Publications 2015.
10. Blackmore R, Gray KM, Melvin GA, Newman L, Boyle JA, Gibson-Helm M. Identifying post-traumatic stress disorder in women of refugee background at a public antenatal clinic. *Arch Womens Ment Health* 2022; 1-8. doi: 10.1007/s00737-021-01167-8
11. Casey LM, Wright M-A, Clough BA. Comparison of perceived barriers and treatment preferences associated with internet-based and face-to-face psychological treatment of depression. *IJCBL* 2014; 4(4):16-22. doi: 10.4018/ijcbl.2014100102
12. Coyne JC, Whiffen VE. Issues in personality as diathesis for depression: the case of sociotropy-dependency and autonomy-self-criticism. *Psychol bull* 1995; 118(3):358. doi: 10.1037/0033-2909.118.3.358
13. Crego C, Widiger TA. Personality disorders. *Psychopathology: Routledge*; 2019; 281-303.
14. Edmonds M, Hadjistavropoulos H, Schneider L, Dear B, Titov N. Who benefits most from therapist-assisted internet-delivered cognitive behaviour therapy in clinical practice? Predictors of symptom change and dropout. *J Anxiety Disord* 2018; 54:24-32. doi: 10.1016/j.janxdis.2018.01.003
15. Eskedal GA, Demetri JM. Etiology and treatment of cluster C personality disorders. *J Ment Health Couns* 2006; 28(1):1-17. doi: 10.17744/mehc.28.1.7yjq6tabcexb1a6j
16. Faith C. Dependent personality disorder: A review of etiology and treatment. *GJCP* 2009; 1(2):7.
17. Fath N, Azad-Fallah P, Tabatabaei R-z, Kazem S, Rahimi C. Validity and reliability of the inventory of interpersonal problems (IIP-32). *J Clinical psychol* 2013; 5(3):69-80.
18. First MB, Williams JB, Benjamin LS, Spitzer RL. Structured clinical interview for DSM-5 personality disorders: SCID-5-PD: American Psychiatric Association Publishing Washington; 2016.
19. First MB, Williams JB, Karg RS, Spitzer RL. Structured clinical interview for DSM-5-Research version (SCID-5 for DSM-5, research version; SCID-5-RV). Arlington, VA: American Psychiatric Association 2015; 2015:1-94.
20. Fjermestad-Noll J, Ronningstam E, Bach BS, Rosenbaum B, Simonsen E. Perfectionism, shame, and aggression in depressive patients with narcissistic personality disorder. *J Pers Disord* 2020; 34(Supplement):25-41. doi: 10.1521/pedi.2020.34.supp.25
21. Ghaderi D, Mostafaei A, Bayazidi S, Shahnazari M. Investigating the prevalence of personality disorders and its relationship with personality traits among students. *Int J Med Res Health Sci* 2016; 5(4):119-26.
22. Gharrae B, Masoumian S, Zamirinejad S, Yaghmaeezadeh H. Psychometric Properties of the Persian Version of the Self-report Personality Questionnaire for Personality Disorders of DSM-5 (SCID-5-SPQ) in Clinical Samples. *Iran J Psychiatry Clin Psychol* 2022; 27(4):508-19. doi: 10.32598/ijpcp.27.4.2497.3
23. Gharraee B, Shabani A, Masoumian S, Zamirinejad S, Yaghmaeezadeh H, Khanjani S, Ghahremani S. Psychometric Properties of the Persian Structured Clinical Interview for Personality Disorders. *East Asian Arch Psychiatry* 2022; 32(4):95-99. doi: 10.12809/eaap2208
24. Gómez-Penedo JM, Fernández-Álvarez J, Maristany M, Freiberg-Hoffmann A. Inventory of Interpersonal Problems-32 (IIP-32): Psychometric properties and normative data in a clinical sample from Argentina. *Rev Eval* 2022; 22(2):64-75.
25. Gude T, Hoffart A, Hedley L, Rø Ø. The dimensionality of dependent personality disorder. *J Pers Disord* 2004; 18(6):604-10. doi: 10.1521/pedi.18.6.604.54793
26. Hedman E, Ljótsson B, Lindefors N. Cognitive behavior therapy via the Internet: a systematic review of applications,

- clinical efficacy and cost-effectiveness. *Expert Rev Pharmacoeconomics Outcomes Res* 2012; 12(6):745-64. doi: 10.1586/erp.12.67
27. Henry JD, Crawford JR. The short-form version of the Depression Anxiety Stress Scales (DASS-21): Construct validity and normative data in a large non-clinical sample. *Br J Clin Psychol* 2005; 44(2):227-39. doi: 10.1348/014466505X29657
 28. Jacob GA, Hauer A, Köhne S, Assmann N, Schaich A, Schweiger U, Fassbinder E. A schema therapy-based eHealth program for patients with borderline personality disorder (prioivi): naturalistic single-arm observational study. *JMIR Mental Health* 2018; 5(4):e10983. doi: 10.2196/10983
 29. Jiang Y, Wei J, Fritzsche K, Toussaint AC, Li T, Cao J, Zhang L, Zhang Y, Chen H, Wu H, Ma X, Li W, Ren J, Lu W, Leonhart R. Assessment of the structured clinical interview (SCID) for DSM-5 for somatic symptom disorder in general hospital outpatient clinics in China. *BMC psychiatry* 2021; 21(1):144. doi: 10.1186/s12888-021-03126-0
 30. Kemp K, Griffiths J, Campbell S, Lovell K. An exploration of the follow-up up needs of patients with inflammatory bowel disease. *J Crohns Colitis* 2013; 7(9):e386-95. doi: 10.1016/j.crohns.2013.03.001
 31. Kenter RM, van-de-Ven PM, Cuijpers P, Koole G, Niamat S, Gerrits RS, Willems M, Straten AV. Costs and effects of Internet cognitive behavioral treatment blended with face-to-face treatment: Results from a naturalistic study. *Internet Interv* 2015; 2(1):77-83. doi: 10.1016/j.invent.2015.01.001
 32. Kiropoulos LA, Klein B, Austin DW, Gilson K, Pier C, Mitchell J, Ciechomski L. Is internet-based CBT for panic disorder and agoraphobia as effective as face-to-face CBT? *J Anxiety Disord* 2008; 22(8):1273-84. doi: 10.1016/j.janxdis.2008.01.008
 33. Klein JP, Hauer A, Berger T, Fassbinder E, Schweiger U, Jacob G. Protocol for the REVISIT-BPD trial, a randomized controlled trial testing the effectiveness of an internet-based self-management intervention in the treatment of borderline personality disorder (BPD). *Front Psychiatry* 2018; 9:439. doi: 10.3389/fpsy.2018.00439
 34. Klein JP, Hauer-von-Mauschwitz A, Berger T, Fassbinder E, Mayer J, Borgwardt S, Wellhöfer B, Schweiger U, Jacob G. Effectiveness and safety of the adjunctive use of an internet-based self-management intervention for borderline personality disorder in addition to care as usual: results from a randomised controlled trial. *BMJ Open* 2021; 11(9):e047771. doi: 10.1136/bmjopen-2020-047771
 35. Komariah M, Amirah S, Faisal EG, Prayogo SA, Maulana S, Platini H, Suryani S, Yosep I, Arifin H. Efficacy of Internet-Based Cognitive Behavioral Therapy for Depression and Anxiety among Global Population during the COVID-19 Pandemic: A Systematic Review and Meta-Analysis of a Randomized Controlled Trial Study. *Healthcare (Basel)* 2022; 10(7):1224. doi: 10.3390/healthcare10071224
 36. Lappalainen P, Granlund A, Siltanen S, Ahonen S, Vitikainen M, Tolvanen A, Lappalainen R. ACT Internet-based vs face-to-face? A randomized controlled trial of two ways to deliver Acceptance and Commitment Therapy for depressive symptoms: An 18-month follow-up. *Behav Res Ther* 2014; 61:43-54. doi: 10.1016/j.brat.2014.07.006
 37. Linehan MM. Skills training manual for treating borderline personality disorder: Guilford press; 1993.
 38. Mahoney AE, Haskelberg H, Mason E, Millard M, Newby J. Comorbid personality difficulties are not associated with poorer outcomes for online cognitive behaviour therapy for symptoms of anxiety and depression. *Personal Ment Health* 2021; 15(3):173-85. doi: 10.1002/pmh.1506
 39. Matusiewicz AK, Hopwood CJ, Banducci AN, Lejuez C. The effectiveness of cognitive behavioral therapy for personality disorders. *Psychiatr Clin* 2010; 33(3):657-85. doi: 10.1016/j.psyc.2010.04.007
 40. Mohammadkhani P, Forouzan AS, Hooshyari Z, Abasi I. Psychometric properties of Persian version of structured clinical interview for DSM-5-research version (SCID-5-RV): a diagnostic accuracy study. *Iran J Psychiatry Behav Sci* 2020; 14(2). doi: 10.5812/ijpbs.100930
 41. Muran JC, Safran JD, Eubanks CF, Gorman BS. The effect of alliance-focused training on a cognitive-behavioral therapy for personality disorders. *J Consult Clin Psychol* 2018; 86(4):384. doi: 10.1037/ccp0000284
 42. Overholser JC. The dependent personality and interpersonal problems. *J Nerv Ment Dis* 1996; 184(1):8-16. doi: 10.1097/0005053-199601000-00003
 43. Kaiser J, Hanschmidt F, Kersting A. The association between therapeutic alliance and outcome in internet-based psychological interventions: a meta-analysis. *Comput Hum Behav* 2021; 114:106512. doi: 10.1016/j.chb.2020.106512
 44. Rafaeli E. Cognitive-behavioral therapies for personality disorders. *Isr J Psychiatry Relat Sci* 2009; 46(4):290.
 45. Rizvi SL, Dimeff LA, Skutch J, Carroll D, Linehan MM. A pilot study of the DBT coach: an interactive mobile phone application for individuals with borderline personality disorder and substance use disorder. *Behav Ther* 2011; 42(4):589-600. doi: 10.1016/j.beth.2011.01.003
 46. Rost KM, Akins RN, Brown FW, Smith GR. The comorbidity of DSM-III-R personality disorders in somatization disorder. *Gen Hosp Psychiatry* 1992; 14(5):322-6. doi: 10.1016/0163-8343(92)90066

47. Sachse R, Breil J, Sachse M, Fasbender J. Persönlichkeitsstörungen: Leitfaden für eine psychologische Psychotherapie. Göttingen: Hogrefe, 2. Auflage 2013.
48. Somma A, Borroni S, Gialdi G, Carlotta D, Emanuela Giarolli L, Barranca M, Cerioli C, Franzoni C, Masci E, Manini R, Luca Busso S, Ruotolo G, Krueger RF, Markon KE, Fossati A. The Inter-Rater Reliability and Validity of the Italian Translation of the Structured Clinical Interview for DSM-5 Alternative Model for Personality Disorders Module I and Module II: A Preliminary Report on Consecutively Admitted Psychotherapy Outpatients. *J Pers Disord* 2020; 34(Suppl C):95-123. doi: 10.1521/pedi_2020_34_511
49. Spence SH, Donovan C, Graham P. Interpersonal problems. *The Handbook of Clinical Adult Psychology* 1994; 229.
50. Sperry L, Sperry J. Cognitive behavior therapy of DSM-5 personality disorders: Assessment, case conceptualization, and treatment: Routledge; 2015. doi: 10.4324/9780203764084
51. Sperry L. Handbook of diagnosis and treatment of DSM-IV personality disorders: Routledge; 2013.
52. Tisdale MJ, Pendleton L, Marler M. MCMI Characteristics of DSM-II-R Bulimics. *J Pers Assess* 1990; 55(3-4):477-83. doi: 10.1080/00223891.1990.9674085
53. Tisdelle DA, Lawrence JSS. Interpersonal problem-solving competency: Review and critique of the literature. *Clin Psychol Rev* 1986; 6(4):337-56. doi: 10.1016/0272-7358(86)90005-X
54. Trull TJ, Jahng S, Tomko RL, Wood PK, Sher KJ. Revised NESARC personality disorder diagnoses: gender, prevalence, and comorbidity with substance dependence disorders. *J pers Disord* 2010; 24(4):412-26. doi: 10.1521/pedi.2010.24.4.412
55. Van-der-Boom B, Boumparis N, Donker T, de Beurs D, Arntz A, Riper H. Internet-delivered interventions for personality disorders-A scoping review. *Internet Interv* 2022; 100525. doi: 10.1016/j.invent.2022.100525
56. Wagner B, Horn AB, Maercker A. Internet-based versus face-to-face cognitive-behavioral intervention for depression: a randomized controlled non-inferiority trial. *J Affect Disord* 2014; 152:113-21. doi: 10.1016/j.jad.2013.06.032
57. Webb CA, Rosso IM, Rauch SL. Internet-based cognitive behavioral therapy for depression: current progress & future directions. *Harv Rev Psychiatry* 2017; 25(3):114. doi: 10.1097/HRP.0000000000000139
58. Weineland S, Ribbegårdh R, Kivi M, Bygdell A, Larsson A, Vernmark K, Lilja JL. Transitioning from face-to-face treatment to iCBT for youths in primary care-therapists' attitudes and experiences. *Internet Interv* 2020; 22:100356. doi: 10.1016/j.invent.2020.100356
59. Zanin CR, Valerio NI. Cognitive behavior therapy for dependent personality disorder: a case report. *Revista Brasileira de Terapia Comportamental e Cognitiva* 2004; 6(1):81-92. doi: 10.31505/rbtcc.v6i1.68